

AMENDMENTS TO THE CLAIMS

This listing of claims will replace all prior versions of claims in the application.

1. (Currently Amended) A stable meloxicam nanoparticulate composition comprising:
 - (a) particles of meloxicam or a salt thereof having an effective average particle size of less than about 2000 nm; and
 - (b) at least one surface stabilizer adsorbed on the surface of the particles,wherein ~~the meloxicam particles have an effective average particle size of less than about 2000 nm~~ in comparative pharmacokinetic testing with a non-nanoparticulate formulation of meloxicam having the same dosage strength and form, the composition exhibits a smaller T_{max} when compared to a T_{max} of the non-nanoparticulate meloxicam formulation.
2. (Previously Presented) The composition of claim 1, wherein the meloxicam is selected from the group consisting of a crystalline phase, an amorphous phase, a semi-crystalline phase, a semi-amorphous phase, and mixtures thereof.
3. (Previously Presented) The composition of claim 1, wherein the effective average particle size of the nanoparticulate meloxicam particles is selected from the group consisting of less than about 1500 nm, less than about 1000 nm, less than about 900 nm, less than about 800 nm, less than about 700 nm, less than about 600 nm, less than about 500 nm, less than about 400 nm, less than about 300 nm, less than about 250 nm, less than about 200 nm, less than about 100 nm, less than about 75 nm, and less than about 50 nm.
4. (Previously Presented) The composition of claim 1, wherein the composition is formulated for administration selected from the group consisting of oral, pulmonary, rectal, ophthalmic, colonic, parenteral, intracisternal, intravaginal, intraperitoneal, local, buccal, nasal, and topical administration.

5. (Previously Presented) The composition of claim 1, wherein the composition is formulated into a dosage form selected from the group consisting of liquid dispersions, gels, aerosols, ointments, creams, controlled release formulations, fast melt formulations, lyophilized formulations, tablets, capsules, delayed release formulations, extended release formulations, pulsatile release formulations, and mixed immediate release and controlled release formulations.

6. (Previously Presented) The composition of claim 1, wherein the composition further comprises one or more pharmaceutically acceptable excipients, carriers, or a combination thereof.

7. (Previously Presented) The composition of claim 1, wherein the meloxicam is present in an amount selected from the group consisting of from about 99.5% to about 0.001%, from about 95% to about 0.1%, and from about 90% to about 0.5%, by weight, based on the total combined weight of the meloxicam and at least one surface stabilizer, not including other excipients.

8. (Previously Presented) The composition of claim 1, wherein the at least one surface stabilizer is present in an amount selected from the group consisting of from about 0.01% to about 99.5% by weight, from about 0.1% to about 95% by weight, and from about 0.5% to about 90% by weight, based on the total combined dry weight of meloxicam and at least one surface stabilizer, not including other excipients.

9. (Previously Presented) The composition of claim 1, comprising at least one primary surface stabilizer and at least one secondary surface stabilizer.

10. (Previously Presented) The composition of claim 1, wherein the surface stabilizer is selected from the group consisting of an anionic surface stabilizer, a cationic surface stabilizer, and an ionic surface stabilizer.

11. (Previously Presented) The composition of claim 10, wherein at least one surface stabilizer is selected from the group consisting of cetyl pyridinium chloride, gelatin, casein,

phosphatides, dextran, glycerol, gum acacia, cholesterol, tragacanth, stearic acid, benzalkonium chloride, calcium stearate, glycerol monostearate, cetostearyl alcohol, cetomacrogol emulsifying wax, sorbitan esters, polyoxyethylene alkyl ethers, polyoxyethylene castor oil derivatives, polyoxyethylene sorbitan fatty acid esters, polyethylene glycols, dodecyl trimethyl ammonium bromide, polyoxyethylene stearates, colloidal silicon dioxide, phosphates, sodium dodecylsulfate, carboxymethylcellulose calcium, hydroxypropyl celluloses, hydroxypropyl methylcellulose, carboxymethylcellulose sodium, methylcellulose, hydroxyethylcellulose, hydroxypropylmethylcellulose phthalate, noncrystalline cellulose, magnesium aluminum silicate, triethanolamine, polyvinyl alcohol, polyvinylpyrrolidone, 4-(1,1,3,3-tetramethylbutyl)-phenol polymer with ethylene oxide and formaldehyde, poloxamers; poloxamines, a charged phospholipid, dioctylsulfosuccinate, dialkylesters of sodium sulfosuccinic acid, sodium lauryl sulfate, alkyl aryl polyether sulfonates, mixtures of sucrose stearate and sucrose distearate, $C_{18}H_{37}CH_2C(O)N(CH_3)-CH_2(CHOH)_4(CH_2OH)_2$, p-isononylphenoxypoly-(glycidol), decanoyl-N-methylglucamide; n-decyl β -D-glucopyranoside; n-decyl β -D-maltopyranoside; n-dodecyl β -D-glucopyranoside; n-dodecyl β -D-maltoside; heptanoyl-N-methylglucamide; n-heptyl- β -D-glucopyranoside; n-heptyl β -D-thioglucoside; n-hexyl β -D-glucopyranoside; nonanoyl-N-methylglucamide; n-nonyl β -D-glucopyranoside; octanoyl-N-methylglucamide; n-octyl- β -D-glucopyranoside; octyl β -D-thioglucopyranoside; lysozyme, PEG-phospholipid, PEG-cholesterol, PEG-cholesterol derivative, PEG-vitamin A, PEG-vitamin E, and random copolymers of vinyl acetate and vinyl pyrrolidone.

12. (Previously Presented) The composition of claim 10, wherein at least one cationic surface stabilizer is selected from the group consisting of a polymer, a biopolymer, a polysaccharide, a cellulosic, an alginate, a nonpolymeric compound, and a phospholipid.

13. (Currently Amended) The composition of claim 10, wherein the surface stabilizer is selected from the group consisting of cationic lipids, polymethylmethacrylate trimethylammonium bromide, sulfonium compounds, polyvinylpyrrolidone-2-dimethylaminoethyl methacrylate dimethyl sulfate, hexadecyltrimethyl ammonium bromide,

phosphonium compounds, quarternary ammonium compounds, benzyl-di(2-chloroethyl)ethylammonium bromide, coconut trimethyl ammonium chloride, coconut trimethyl ammonium bromide, coconut methyl dihydroxyethyl ammonium chloride, coconut methyl dihydroxyethyl ammonium bromide, decyl triethyl ammonium chloride, decyl dimethyl hydroxyethyl ammonium chloride, decyl dimethyl hydroxyethyl ammonium chloride bromide, C₁₂₋₁₅dimethyl hydroxyethyl ammonium chloride, C₁₂₋₁₅dimethyl hydroxyethyl ammonium chloride bromide, coconut dimethyl hydroxyethyl ammonium chloride, coconut dimethyl hydroxyethyl ammonium bromide, myristyl trimethyl ammonium methyl sulphate, lauryl dimethyl benzyl ammonium chloride, lauryl dimethyl benzyl ammonium bromide, lauryl dimethyl (ethenoxy)₄ ammonium chloride, lauryl dimethyl (ethenoxy)₄ ammonium bromide, N-alkyl (C₁₂₋₁₈)dimethylbenzyl ammonium chloride, N-alkyl (C₁₄₋₁₈)dimethyl-benzyl ammonium chloride, N-tetradecyldimethylbenzyl ammonium chloride monohydrate, dimethyl didecyl ammonium chloride, N-alkyl and (C₁₂₋₁₄) dimethyl 1-naphthylmethyl ammonium chloride, trimethylammonium halide, alkyl-trimethylammonium salts, dialkyl-dimethylammonium salts, lauryl trimethyl ammonium chloride, ethoxylated alkyamidoalkyldialkylammonium salt, an ethoxylated trialkyl ammonium salt, dialkylbenzene dialkylammonium chloride, N-didecyldimethyl ammonium chloride, N-tetradecyldimethylbenzyl ammonium, chloride monohydrate, N-alkyl(C₁₂₋₁₄) dimethyl 1-naphthylmethyl ammonium chloride, dodecyldimethylbenzyl ammonium chloride, dialkyl benzenealkyl ammonium chloride, lauryl trimethyl ammonium chloride, alkylbenzyl methyl ammonium chloride, alkyl benzyl dimethyl ammonium bromide, C₁₂ trimethyl ammonium bromides, C₁₅ trimethyl ammonium bromides, C₁₇ trimethyl ammonium bromides, dodecylbenzyl triethyl ammonium chloride, poly-diallyldimethylammonium chloride (DADMAC), dimethyl ammonium chlorides, alkyl dimethylammonium halogenides, tricetyl methyl ammonium chloride, decyltrimethylammonium bromide, dodecyltriethylammonium bromide, tetradecyltrimethylammonium bromide, methyl trioctylammonium chloride, ~~POLYQUAT 10™~~ polyquaternium 10, tetrabutylammonium bromide, benzyl trimethylammonium bromide, choline esters, benzalkonium chloride, stearalkonium chloride compounds, cetyl pyridinium bromide,

cetyl pyridinium chloride, halide salts of quaternized polyoxyethylalkylamines, MIRAPOL™, ALKAQUAT™ quaternized ammonium salt polymers, alkyl pyridinium salts; amines, amine salts, amine oxides, imide azolinium salts, protonated quaternary acrylamides, methylated quaternary polymers, and cationic guar.

14. (Previously Presented) The composition of claim 1, wherein the T_{\max} of the composition, when assayed in the plasma of the mammalian subject, is selected from the group consisting of less than about 5 hours, less than about 4 hours, less than about 3 hours, less than about 2 hours, less than about 1 hour, less than less than about 50 minutes, less than about 45 minutes, less than about 40 minutes, less than about 35 minutes, less than about 30 minutes, less than about 20 minutes, less than about 15 minutes, less than about 10 minutes, and less than about 5 minutes.

15. (Currently Amended) The composition of claim 1, wherein ~~in comparative pharmacokinetic testing with a standard non-nanoparticulate commercial formulation of meloxicam, the composition of claim 1 exhibits a~~ the T_{\max} of the composition is selected from the group consisting of less than about 90%, less than about 80%, less than about 70%, less than about 50%, less than about 30%, less than about 25%, less than about 20%, less than about 15%, and less than about 10% of the T_{\max} exhibited by the ~~standard commercial non-nanoparticulate meloxicam formulation.~~

16. (Previously Presented) The composition of claim 1, wherein the C_{\max} of the composition, when assayed in the plasma of the mammalian subject, is selected from the group consisting of greater than about 1 $\mu\text{g/mL}$, greater than about 3 $\mu\text{g/mL}$, greater than about 5 $\mu\text{g/mL}$, greater than about 10 $\mu\text{g/mL}$, and greater than about 15 $\mu\text{g/mL}$.

17. (Currently Amended) The composition of claim 1, wherein ~~in comparative pharmacokinetic testing with a standard non-nanoparticulate commercial formulation of meloxicam, the composition of claim 1 exhibits a~~ the C_{\max} of the composition is selected from the group consisting of greater than about 20%, greater than about 40%, greater than about 60%,

greater than about 80%, greater than about 100%, greater than about 140%, greater than about 180%, greater than about 200%, greater than about 240%, greater than about 280%, greater than about 300%, greater than about 340%, greater than about 380%, and greater than about 400% of the C_{max} exhibited by a ~~standard commercial~~ non-nanoparticulate meloxicam formulation.

18. (Previously Presented) The composition of claim 1, additionally comprising a meloxicam composition having an effective average particle size which is greater than about 2 microns.

19. (Previously Presented) The composition of claim 1, additionally comprising at least one additional nanoparticulate meloxicam composition, having an effective average particle size of less than about 2 microns, wherein said additional nanoparticulate meloxicam composition has an effective average particle size which is different than particle size of the nanoparticulate meloxicam composition of claim 1.

20. (Previously Presented) The composition of claim 1, additionally comprising at least one non-meloxicam active agent selected from the group consisting of proteins, peptides, nucleotides, anti-obesity drugs, nutraceuticals, dietary supplements, carotenoids, corticosteroids, elastase inhibitors, anti-fungals, alkylxanthine, oncology therapies, anti-emetics, analgesics, opioids, antipyretics, cardiovascular agents, anti-inflammatory agents, anthelmintics, anti-arrhythmic agents, antibiotics, anticoagulants, antidepressants, antidiabetic agents, antiepileptics, antihistamines, antihypertensive agents, antimuscarinic agents, antimycobacterial agents, antineoplastic agents, immunosuppressants, antithyroid agents, antiviral agents, anxiolytics, sedatives, astringents, beta-adrenoceptor blocking agents, blood products, blood substitutes, cardiac inotropic agents, contrast media, corticosteroids, cough suppressants, diagnostic agents, diagnostic imaging agents, diuretics, dopaminergics, haemostatics, immunological agents, lipid regulating agents, muscle relaxants, parasympathomimetics, parathyroid calcitonin and biphosphonates, prostaglandins, radio-pharmaceuticals, sex hormones, anti-allergic agents, stimulants, anoretics, sympathomimetics, thyroid agents, Mu receptor antagonists, Kappa

receptor antagonists, non-narcotic analgesics, monoamine uptake inhibitors, adenosine regulating agents, cannabinoid derivatives, Substance P antagonists, neurokinin-1 receptor antagonists, and sodium channel blockers.

21. (Previously Presented) The composition of claim 20, wherein said nutraceutical is selected from the group consisting of lutein, folic acid, fatty acids, fruit extracts, vegetable extracts, vitamin supplements, mineral supplements, phosphatidylserine, lipoic acid, melatonin, glucosamine/chondroitin, Aloe Vera, Guggul, glutamine, amino acids, green tea, lycopene, whole foods, food additives, herbs, phytonutrients, antioxidants, flavonoid constituents of fruits, evening primrose oil, flax seeds, fish oils, marine animal oils, and probiotics.

22. (Previously Presented) The composition of claim 20, wherein said anti-inflammatory agent is a COX-2 inhibitor selected from the group consisting of celecoxib, rofecoxib, valdecoxib, parecoxib, MK-966, etoricoxib, 4-[5-(4-chlorophenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl]] benzenesulfonamide, N-(2-cyclohexyloxy-4-nitrophenyl)methane sulfonamide, methyl sulfone spiro(2.4)hept-5-ene I, SC-57666, celecoxib, SC-558, SC-560, etodolac, 5,5-dimethyl-3-(3-fluorophenyl)-4-(4-methylsulfonyl)phenyl 2(5H)-furanone, MK-476, L-745337, L-761066, L-761000, L-748780, L-748731, 5-Bromo-2-(4-fluorophenyl)-3-(4-(methylsulfonyl)phenyl), 1-(7-tert.-butyl-2,3-dihydro-3,3-dimethylbenzo(b)furan-5-yl)-4-cyclopropylbutan-1-one, 3-formylamino-7-methylsulfonylamino-6-phenoxy-4H-1-benzopyran-4-one, BF 389, PD 136005, PD 142893, PD 145065, flurbiprofen, nimesulide, nabumetone, flosulide, piroxicam, dicofenac, COX-189, D 1367, 4 nitro 2 phenoxymethane sulfonanilide, (3 benzoyldifluoromethane sulfonanilide, diflumidone), JTE-522, 4'-Acetyl-2'-(2,4-difluorophenoxy)methanesulfonanilide, FK 867, FR 115068, GR 253035, RWJ 63556, RWJ 20485, ZK 38997, (E)-(5)-(3,5-di-tert-butyl-4-hydroxybenzylidene)-2-ethyl-1,2-isothiazolidine-1,1-dioxide indomethacin, CL 1004, RS 57067, RS 104894, SC 41930, SB 205312, SKB 209670, and Ono 1078.

23. (Previously Presented) The composition of claim 20, wherein said non-meloxicam active agent is selected from the group consisting of aceclofenac, acemetacin, e-acetamidocaproic acid, acetaminophen, acetaminosalol, acetanilide, acetylsalicylic acid, S-adenosylmethionine, alclofenac, alfentanil, allylprodine, alminoprofen, aloxiprin, alphaproline, aluminum bis(acetylsalicylate), amfenac, aminochlorthenoxazin, 3-amino-4-hydroxybutyric acid, 2-amino-4-picoline, aminopropyl, aminopyrine, amixetrine, ammonium salicylate, ampiroxicam, amtolmetin guacil, anileridine, antipyrine, antipyrine salicylate, antrafenine, apazone, bendazac, benorylate, benoxaprofen, benzpiperylon, benzydamine, benzylmorphine, bermoprofen, bezitramide, α -bisabolol, bromfenac, p-bromoacetanilide, 5-bromosalicylic acid acetate, bromosaligenin, bucetin, bucloxic acid, bucolome, bufexamac, bumadizon, buprenorphine, butacetin, butibufen, butophanol, calcium acetylsalicylate, carbamazepine, carbiphene, carprofen, carsalam, chlorobutanol, chlorthenoxazin, choline salicylate, cinchophen, cinmetacin, ciramadol, clidanac, clometacin, clonitazene, clonixin, clopirac, clove, codeine, codeine methyl bromide, codeine phosphate, codeine sulfate, cropropamide, crotethamide, desomorphine, dexoxadrol, dextromoramide, dezocine, diampromide, diclofenac sodium, difenamizole, difenpiramide, diflunisal, dihydrocodeine, dihydrocodeinone enol acetate, dihydromorphine, dihydroxyaluminum acetylsalicylate, dimenoxadol, dimepheptanol, dimethylthiambutene, dioxaphetyl butyrate, dipipanone, diprocetyl, dipyrone, ditazol, droxicam, emorfazone, enfenamic acid, epirizole, eptazocine, etersalate, ethenzamide, ethoheptazine, ethoxazene, ethylmethylthiambutene, ethylmorphine, etodolac, etofenamate, etonitazene, eugenol, felbinac, fenbufen, fenclozic acid, fendosal, fenoprofen, fentanyl, fentiazac, fepradinol, feprazone, floctafenine, flufenamic acid, flunoxaprofen, fluoresone, flupirtine, fluproquazone, flurbiprofen, fosfosal, gentisic acid, glafenine, glucametacin, glycol salicylate, guaiazulene, hydrocodone, hydromorphone, hydroxypethidine, ibufenac, ibuprofen, ibuproxam, imidazole salicylate, indomethacin, indoprofen, isofezolac, isoladol, isomethadone, isonixin, isoxepac, isoxicam, ketobemidone, ketoprofen, ketorolac, p-lactophenetide, lefetamine, levorphanol, lofentanil, lonazolac, lomoxicam, loxoprofen, lysine acetylsalicylate, magnesium acetylsalicylate, meclofenamic acid, mefenamic acid, meperidine, meptazinol, mesalamine, metazocine,

methadone hydrochloride, methotrimeprazine, metiazinic acid, metofoline, metopon, mofebutazone, mofezolac, morazone, morphine, morphine hydrochloride, morphine sulfate, morpholine salicylate, myrophine, nabumetone, nalbuphine, 1-naphthyl salicylate, naproxen, narceine, nefopam, nicomorphine, nifenazone, niflumic acid, nimesulide, 5'-nitro-2'-propoxyacetanilide, norlevorphanol, normethadone, normorphine, norpipanone, olsalazine, opium, oxaceprol, oxametacine, oxaprozin, oxycodone, oxymorphone, oxyphenbutazone, papaveretum, paranyline, parsalmide, pentazocine, perisoxal, phenacetin, phenadoxone, phenazocine, phenazopyridine hydrochloride, phenocoll, phenoperidine, phenopyrazone, phenyl acetylsalicylate, phenylbutazone, phenyl salicylate, phenylamidol, piketoprofen, piminodine, pipebuzone, piperylone, pirofen, pirazolac, piritramide, piroxicam, pranoprofen, proglumetacin, proheptazine, promedol, propacetamol, propiram, propoxyphene, propyphenazone, proquazone, protizinic acid, ramifenazone, remifentanil, rimazolium metilsulfate, salacetamide, salicin, salicylamide, salicylamide o-acetic acid, salicylsulfuric acid, salsalte, salverine, simetride, sodium salicylate, sufentanil, sulfasalazine, sulindac, superoxide dismutase, suprofen, suxibuzone, talniflumate, tenidap, tenoxicam, terofenamate, tetrandrine, thiazolinobutazone, tiaprofenic acid, tiaramide, tilidine, tinoridine, tolfenamic acid, tolmetin, tramadol, tropesin, viminol, xenbucin, ximoprofen, zaltoprofen, and zomepirac.

24. (Previously Presented) The composition of any of claims 20, 21, 22, or 23, wherein at least one the non-meloxicam active agents has an effective average particle size of less than about 2 microns.

25. (Previously Presented) The composition of any of 20, 21, 22, or 23, wherein at least one the non-meloxicam active agents is a conventional particle sized active agent.

26. (Previously Presented) A method of making a nanoparticulate composition comprising contacting meloxicam particles with at least one surface stabilizer for a time and under conditions sufficient to provide a nanoparticulate meloxicam composition having an effective average particle size of less than about 2000 nm.

27. (Previously Presented) The method of claim 26, wherein said contacting comprises grinding.

28. (Previously Presented) The method of claim 27, wherein said grinding comprises wet grinding.

29. (Previously Presented) The method of claim 26, wherein said contacting comprises homogenizing.

30. (Previously Presented) The method of claim 26, wherein said contacting comprises:

- (a) dissolving the meloxicam particles in a solvent;
- (b) adding the resulting meloxicam solution to a solution comprising at least one surface stabilizer; and
- (c) precipitating the solubilized meloxicam having at least one surface stabilizer associated with the surface thereof by the addition thereto of a non-solvent.

31. (Previously Presented) The method of claim 26, wherein the meloxicam is selected from the group consisting of a crystalline phase, an amorphous phase, a semi-crystalline phase, a semi-amorphous phase, and mixtures thereof.

32. (Previously Presented) The method of claim 26, wherein the effective average particle size of the nanoparticulate meloxicam particles is selected from the group consisting of less than about 1500 nm, less than about 1000 nm, less than about 900 nm, less than about 800 nm, less than about 700 nm, less than about 600 nm, less than about 500 nm, less than about 400 nm, less than about 300 nm, less than about 250 nm, less than about 200 nm, less than about 100 nm, less than about 75 nm, and less than about 50 nm.

33. (Previously Presented) The method of claim 26, wherein the composition is formulated for an administration form selected from the group consisting of oral, pulmonary,

rectal, ophthalmic, colonic, parenteral, intracisternal, intravaginal, intraperitoneal, local, buccal, nasal, and topical administration.

34. (Previously Presented) The method of claim 26, wherein the composition is formulated into a dosage form selected from the group consisting of liquid dispersions, gels, aerosols, ointments, creams, controlled release formulations, fast melt formulations, lyophilized formulations, tablets, capsules, delayed release formulations, extended release formulations, pulsatile release formulations, and mixed immediate release and controlled release formulations.

35. (Previously Presented) The method of claim 26, wherein the composition further comprises one or more pharmaceutically acceptable excipients, carriers, or a combination thereof.

36. (Previously Presented) The method of claim 26, wherein the meloxicam is present in an amount selected from the group consisting of from about 99% to about 0.001%, from about 95% to about 0.5%, and from about 90% to about 0.5%, by weight, based on the total combined weight of the meloxicam and at least one surface stabilizer, not including other excipients.

37. (Previously Presented) The method of claim 26, wherein at least one surface stabilizer is present in an amount selected from the group consisting of from about 0.01% to about 99.5% by weight, from about 0.1% to about 95% by weight, and from about 0.5% to about 90% by weight, based on the total combined dry weight of the meloxicam and at least one surface stabilizer, not including other excipients.

38. (Previously Presented) The method of claim 26, comprising at least one primary surface stabilizer and at least one secondary surface stabilizer.

39. (Previously Presented) The method of claim 26, wherein the surface stabilizer is selected from the group consisting of an anionic surface stabilizer, a cationic surface stabilizer, and an ionic surface stabilizer.

40. (Previously Presented) The method of claim 39, wherein at least one surface stabilizer is selected from the group consisting of cetyl pyridinium chloride, gelatin, casein, phosphatides, dextran, glycerol, gum acacia, cholesterol, tragacanth, stearic acid, benzalkonium chloride, calcium stearate, glycerol monostearate, cetostearyl alcohol, cetomacrogol emulsifying wax, sorbitan esters, polyoxyethylene alkyl ethers, polyoxyethylene castor oil derivatives, polyoxyethylene sorbitan fatty acid esters, polyethylene glycols, dodecyl trimethyl ammonium bromide, polyoxyethylene stearates, colloidal silicon dioxide, phosphates, sodium dodecylsulfate, carboxymethylcellulose calcium, hydroxypropyl celluloses, hydroxypropyl methylcellulose, carboxymethylcellulose sodium, methylcellulose, hydroxyethylcellulose, hydroxypropylmethylcellulose phthalate, noncrystalline cellulose, magnesium aluminum silicate, triethanolamine, polyvinyl alcohol, polyvinylpyrrolidone, 4-(1,1,3,3-tetramethylbutyl)-phenol polymer with ethylene oxide and formaldehyde, poloxamers; poloxamines, a charged phospholipid, dioctylsulfosuccinate, dialkylesters of sodium sulfosuccinic acid, sodium lauryl sulfate, alkyl aryl polyether sulfonates, mixtures of sucrose stearate and sucrose distearate, $C_{18}H_{37}CH_2C(O)N(CH_3)-CH_2(CHOH)_4(CH_2OH)_2$, p-isononylphenoxypoly-(glycidol), decanoyl-N-methylglucamide; n-decyl β -D-glucopyranoside; n-decyl β -D-maltopyranoside; n-dodecyl β -D-glucopyranoside; n-dodecyl β -D-maltoside; heptanoyl-N-methylglucamide; n-heptyl- β -D-glucopyranoside; n-heptyl β -D-thioglucoside; n-hexyl β -D-glucopyranoside; nonanoyl-N-methylglucamide; n-nonyl β -D-glucopyranoside; octanoyl-N-methylglucamide; n-octyl- β -D-glucopyranoside; octyl β -D-thioglucopyranoside; lysozyme, PEG-phospholipid, PEG-cholesterol, PEG-cholesterol derivative, PEG-vitamin A, PEG-vitamin E, and random copolymers of vinyl acetate and vinyl pyrrolidone.

41. (Previously Presented) The method of claim 39, wherein at least one cationic surface stabilizer is selected from the group consisting of a polymer, a biopolymer, a polysaccharide, a cellulosic, an alginate, a nonpolymeric compound, and a phospholipid.

42. (Currently Amended) The method of claim 39, wherein the surface stabilizer is selected from the group consisting of cationic lipids, polymethylmethacrylate

trimethylammonium bromide, sulfonium compounds, polyvinylpyrrolidone-2-dimethylaminoethyl methacrylate dimethyl sulfate, hexadecyltrimethyl ammonium bromide, phosphonium compounds, quaternary ammonium compounds, benzyl-di(2-chloroethyl)ethylammonium bromide, coconut trimethyl ammonium chloride, coconut trimethyl ammonium bromide, coconut methyl dihydroxyethyl ammonium chloride, coconut methyl dihydroxyethyl ammonium bromide, decyl triethyl ammonium chloride, decyl dimethyl hydroxyethyl ammonium chloride, decyl dimethyl hydroxyethyl ammonium chloride bromide, C₁₂₋₁₅dimethyl hydroxyethyl ammonium chloride, C₁₂₋₁₅dimethyl hydroxyethyl ammonium chloride bromide, coconut dimethyl hydroxyethyl ammonium chloride, coconut dimethyl hydroxyethyl ammonium bromide, myristyl trimethyl ammonium methyl sulphate, lauryl dimethyl benzyl ammonium chloride, lauryl dimethyl benzyl ammonium bromide, lauryl dimethyl (ethenoxy)₄ ammonium chloride, lauryl dimethyl (ethenoxy)₄ ammonium bromide, N-alkyl (C₁₂₋₁₈)dimethylbenzyl ammonium chloride, N-alkyl (C₁₄₋₁₈)dimethyl-benzyl ammonium chloride, N-tetradecyldimethylbenzyl ammonium chloride monohydrate, dimethyl didecyl ammonium chloride, N-alkyl and (C₁₂₋₁₄) dimethyl 1-naphthylmethyl ammonium chloride, trimethylammonium halide, alkyl-trimethylammonium salts, dialkyl-dimethylammonium salts, lauryl trimethyl ammonium chloride, ethoxylated alkyamidoalkyldialkylammonium salt, an ethoxylated trialkyl ammonium salt, dialkylbenzene dialkylammonium chloride, N-didecyldimethyl ammonium chloride, N-tetradecyldimethylbenzyl ammonium, chloride monohydrate, N-alkyl(C₁₂₋₁₄) dimethyl 1-naphthylmethyl ammonium chloride, dodecyldimethylbenzyl ammonium chloride, dialkyl benzenealkyl ammonium chloride, lauryl trimethyl ammonium chloride, alkylbenzyl methyl ammonium chloride, alkyl benzyl dimethyl ammonium bromide, C₁₂ trimethyl ammonium bromides, C₁₅ trimethyl ammonium bromides, C₁₇ trimethyl ammonium bromides, dodecylbenzyl triethyl ammonium chloride, polydiallyldimethylammonium chloride (DADMAC), dimethyl ammonium chlorides, alkyldimethylammonium halogenides, tricetyl methyl ammonium chloride, decyltrimethylammonium bromide, dodecyltriethylammonium bromide, tetradecyltrimethylammonium bromide, methyl trioctylammonium chloride, ~~POLYQUAT 10™~~

polyquaternium 10, tetrabutylammonium bromide, benzyl trimethylammonium bromide, choline esters, benzalkonium chloride, stearylalkonium chloride compounds, cetyl pyridinium bromide, cetyl pyridinium chloride, halide salts of quaternized polyoxyethylalkylamines, ~~MIRAPOL™~~, ~~ALKAQUAT™~~ quaternized ammonium salt polymers, alkyl pyridinium salts; amines, amine salts, amine oxides, imide azolinium salts, protonated quaternary acrylamides, methylated quaternary polymers, and cationic guar.

43. (Previously Presented) The method of claim 26, wherein after preparation of a first nanoparticulate meloxicam composition, a second meloxicam composition having an effective average particle size of greater than about 2 microns is combined with the first nanoparticulate meloxicam composition.

44. (Previously Presented) The method of claim 26, wherein either prior to or subsequent to preparation of the nanoparticulate meloxicam composition, at least one non-meloxicam active agent is added to the meloxicam composition, wherein said non-meloxicam active agent is selected from the group consisting of proteins, peptides, nucleotides, anti-obesity drugs, nutraceuticals, dietary supplements, carotenoids, corticosteroids, elastase inhibitors, anti-fungals, alkylxanthine, oncology therapies, anti-emetics, analgesics, opioids, antipyretics, cardiovascular agents, anti-inflammatory agents, anthelmintics, anti-arrhythmic agents, antibiotics, anticoagulants, antidepressants, antidiabetic agents, antiepileptics, antihistamines, antihypertensive agents, antimuscarinic agents, antimycobacterial agents, antineoplastic agents, immunosuppressants, antithyroid agents, antiviral agents, anxiolytics, sedatives, astringents, beta-adrenoceptor blocking agents, blood products, blood substitutes, cardiac inotropic agents, contrast media, corticosteroids, cough suppressants, diagnostic agents, diagnostic imaging agents, diuretics, dopaminergics, haemostatics, immunological agents, lipid regulating agents, muscle relaxants, parasympathomimetics, parathyroid calcitonin and biphosphonates, prostaglandins, radio- pharmaceuticals, sex hormones, anti-allergic agents, stimulants, anoretics, sympathomimetics, thyroid agents, Mu receptor antagonists, Kappa receptor antagonists, non-narcotic analgesics, monoamine uptake inhibitors, adenosine regulating agents, cannabinoid

derivatives, Substance P antagonists, neurokinin-1 receptor antagonists, and sodium channel blockers.

45. (Previously Presented) The method of claim 44, wherein said nutraceutical is selected from the group consisting of lutein, folic acid, fatty acids, fruit extracts, vegetable extracts, vitamin supplements, mineral supplements, phosphatidylserine, lipoic acid, melatonin, glucosamine/chondroitin, Aloe Vera, Guggul, glutamine, amino acids, green tea, lycopene, whole foods, food additives, herbs, phytonutrients, antioxidants, flavonoid constituents of fruits, evening primrose oil, flax seeds, fish oils, marine animal oils, and probiotics.

46. (Previously Presented) The method of claim 44, wherein said anti-inflammatory agent is a COX-2 inhibitor selected from the group consisting of celecoxib, rofecoxib, valdecoxib, parecoxib, MK-966, etoricoxib, 4-[5-(4-chlorophenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl]] benzenesulfonamide, N-(2-cyclohexyloxy-4-nitrophenyl)methane sulfonamide, methyl sulfone spiro(2.4)hept-5-ene I, SC-57666, celecoxib, SC-558, SC-560, etodolac, 5,5-dimethyl-3-(3-fluorophenyl)-4-(4-methylsulfonyl)phenyl 2(5H)-furanone, MK-476, L-745337, L-761066, L-761000, L-748780, L-748731, 5-Bromo-2-(4-fluorophenyl)-3-(4-(methylsulfonyl)phenyl, 1-(7-tert.-butyl-2,3-dihydro-3,3-dimethylbenzo(b)furan-5-yl)-4-cyclopropylbutan-1-one, 3-formylamino-7-methylsulfonylamino-6-phenoxy-4H-1-benzopyran-4-one, BF 389, PD 136005, PD 142893, PD 145065, flurbiprofen, nimesulide, nabumetone, flosulide, piroxicam, dicofenac, COX-189, D 1367, 4 nitro 2 phenoxymethane sulfonanilide, (3 benzoyldifluoromethane sulfonanilide, diflumidone), JTE-522, 4'-Acetyl-2'-(2,4-difluorophenoxy)methanesulfonanilide, FK 867, FR 115068, GR 253035, RWJ 63556, RWJ 20485, ZK 38997, (E)-(5)-(3,5-di-tert-butyl-4-hydroxybenzylidene)-2-ethyl-1,2-isothiazolidine-1,1-dioxide indomethacin, CL 1004, RS 57067, RS 104894, SC 41930, SB 205312, SKB 209670, and Ono 1078.

47. (Previously Presented) The method of claim 44, wherein said non-meloxicam active agent is selected from the group consisting of aceclofenac, acemetacin, e-acetamidocaproic

acid, acetaminophen, acetaminosalol, acetanilide, acetylsalicylic acid, S-adenosylmethionine, alclofenac, alfentanil, allylprodine, alminoprofen, aloxiprin, alphaprodine, aluminum bis(acetylsalicylate), amfenac, aminochlorthenoxazin, 3-amino-4-hydroxybutyric acid, 2-amino-4-picoline, aminopropylon, aminopyrine, amixetrine, ammonium salicylate, ampiroxicam, amtolmetin guacil, anileridine, antipyrine, antipyrine salicylate, antrafenine, apazone, bendazac, benorylate, benoxaprofen, benzpiperylon, benzydamine, benzylmorphine, bermoprofen, bezitramide, α -bisabolol, bromfenac, p-bromoacetanilide, 5-bromosalicylic acid acetate, bromosaligenin, bucetin, bucloxic acid, bucolome, bufexamac, bumadizon, buprenorphine, butacetin, butibufen, butophanol, calcium acetylsalicylate, carbamazepine, carbiphen, carprofen, carsalam, chlorobutanol, chlorthenoxazin, choline salicylate, cinchophen, cinmetacin, ciramadol, clidanac, clometacin, clonitazene, clonixin, clopirac, clove, codeine, codeine methyl bromide, codeine phosphate, codeine sulfate, cropropamide, crotethamide, desomorphine, dexoxadrol, dextromoramide, dezocine, diampromide, diclofenac sodium, difenamizole, difenpiramide, diflunisal, dihydrocodeine, dihydrocodeinone enol acetate, dihydromorphine, dihydroxyaluminum acetylsalicylate, dimenoxadol, dimepheptanol, dimethylthiambutene, dioxaphetyl butyrate, dipipanone, diprocetyl, dipyrone, ditazol, droxicam, emorfazone, enfenamic acid, epirizole, eptazocine, etersalate, ethenzamide, ethoheptazine, ethoxazene, ethylmethylthiambutene, ethylmorphine, etodolac, etofenamate, etonitazene, eugenol, felbinac, fenbufen, fenclozic acid, fendosal, fenoprofen, fentanyl, fentiazac, fepradinol, feprazone, floctafenine, flufenamic acid, flunoxaprofen, fluoresone, flupirtine, fluproquazone, flurbiprofen, fosfosal, gentisic acid, glafenine, glucametacin, glycol salicylate, guaiazulene, hydrocodone, hydromorphone, hydroxypethidine, ibufenac, ibuprofen, ibuproxam, imidazole salicylate, indomethacin, indoprofen, isofezolac, isoladol, isomethadone, isonixin, isoxepac, isoxicam, ketobemidone, ketoprofen, ketorolac, p-lactophenetide, lefetamine, levorphanol, lofentanil, lonazolac, lomoxicam, loxoprofen, lysine acetylsalicylate, magnesium acetylsalicylate, meclofenamic acid, mefenamic acid, meperidine, meptazinol, mesalamine, metazocine, methadone hydrochloride, methotrimeprazine, metiazinic acid, metofoline, metopon, mofebutazone, mofezolac, morazone, morphine, morphine hydrochloride, morphine sulfate,

morpholine salicylate, myrophine, nabumetone, nalbuphine, 1-naphthyl salicylate, naproxen, narceine, nefopam, nicomorphine, nifenazone, niflumic acid, nimesulide, 5'-nitro-2'-propoxyacetanilide, norlevorphanol, normethadone, normorphine, norpipanone, olsalazine, opium, oxaceprol, oxametacine, oxaprozin, oxycodone, oxymorphone, oxyphenbutazone, papaveretum, paranyline, parsalimide, pentazocine, perisoxal, phenacetin, phenadoxone, phenazocine, phenazopyridine hydrochloride, phenocoll, phenoperidine, phenopyrazone, phenyl acetylsalicylate, phenylbutazone, phenyl salicylate, phenylramidol, piketoprofen, piminodine, pipebuzone, piperylone, piprofen, pirazolac, piritramide, piroxicam, pranoprofen, proglumetacin, proheptazine, promedol, propacetamol, propiram, propoxyphene, propyphenazone, proquazone, protizinic acid, ramifenazone, remifentanil, rimazolium metilsulfate, salacetamide, salicin, salicylamide, salicylamide o-acetic acid, salicylsulfuric acid, salsalte, salverine, simetride, sodium salicylate, sufentanil, sulfasalazine, sulindac, superoxide dismutase, suprofen, suxibuzone, talniflumate, tenidap, tenoxicam, terofenamate, tetrandrine, thiazolinobutazone, tiaprofenic acid, tiaramide, tilidine, tinoridine, tolfenamic acid, tolmetin, tramadol, tropesin, viminol, xenbucin, ximoprofen, zaltoprofen, and zomepirac.

48. (Previously Presented) The method of any of claims 44, 45, 46, or 47, wherein at least one non-meloxicam active agent has an effective average particle size of less than about 2 microns.

49. (Previously Presented) The method of any of claims 44, 45, 46, or 47, wherein at least one non-meloxicam active agent has an effective average particle size of greater than about 2 microns.

50. (Previously Presented) A method of treating a subject in need with a nanoparticulate meloxicam formulation comprising administering to the subject an effective amount of a nanoparticulate composition comprising meloxicam particles and at least one surface stabilizer, wherein the meloxicam particles have an effective average particle size of less than about 2000 nm.

51. (Previously Presented) The method of claim 50, wherein the meloxicam is selected from the group consisting of a crystalline phase, an amorphous phase, a semi-crystalline phase, a semi-amorphous phase, and mixtures thereof.

52. (Previously Presented) The method of claim 50, wherein the effective average particle size of the nanoparticulate meloxicam particles is selected from the group consisting of less than about 1500 nm, less than about 1000 nm, less than about 900 nm, less than about 800 nm, less than about 700 nm, less than about 600 nm, less than about 500 nm, less than about 400 nm, less than about 300 nm, less than about 250 nm, less than about 200 nm, less than about 100 nm, less than about 75 nm, and less than about 50 nm.

53. (Previously Presented) The method of claim 50, wherein the composition is formulated for an administration form selected from the group consisting of oral, pulmonary, rectal, ophthalmic, colonic, parenteral, intracisternal, intravaginal, intraperitoneal, local, buccal, nasal, and topical administration.

54. (Previously Presented) The method of claim 50, wherein the composition is a dosage form selected from the group consisting of liquid dispersions, gels, aerosols, ointments, creams, controlled release formulations, fast melt formulations, lyophilized formulations, tablets, capsules, delayed release formulations, extended release formulations, pulsatile release formulations, and mixed immediate release and controlled release formulations.

55. (Previously Presented) The method of claim 50, wherein the composition further comprises one or more pharmaceutically acceptable excipients, carriers, or a combination thereof.

56. (Previously Presented) The method of claim 50, wherein the meloxicam is present in an amount selected from the group consisting of from about 99% to about 0.001%, from about 95% to about 0.5%, and from about 90% to about 0.5%, by weight, based on the total

combined weight of the meloxicam and at least one surface stabilizer, not including other excipients.

57. (Previously Presented) The method of claim 50, wherein at least one surface stabilizer is present in an amount selected from the group consisting of from about 0.01% to about 99.5% by weight, from about 0.1% to about 95% by weight, and from about 0.5% to about 90% by weight, based on the total combined dry weight of meloxicam and at least one surface stabilizer, not including other excipients.

58. (Previously Presented) The method of claim 50, comprising at least one primary surface stabilizer and at least one secondary surface stabilizer.

59. (Previously Presented) The method of claim 50, wherein the surface stabilizer is selected from the group consisting of an anionic surface stabilizer, a cationic surface stabilizer, and an ionic surface stabilizer.

60. (Previously Presented) The method of claim 59, wherein the at least one surface stabilizer is selected from the group consisting of cetyl pyridinium chloride, gelatin, casein, phosphatides, dextran, glycerol, gum acacia, cholesterol, tragacanth, stearic acid, benzalkonium chloride, calcium stearate, glycerol monostearate, cetostearyl alcohol, cetomacrogol emulsifying wax, sorbitan esters, polyoxyethylene alkyl ethers, polyoxyethylene castor oil derivatives, polyoxyethylene sorbitan fatty acid esters, polyethylene glycols, dodecyl trimethyl ammonium bromide, polyoxyethylene stearates, colloidal silicon dioxide, phosphates, sodium dodecylsulfate, carboxymethylcellulose calcium, hydroxypropyl celluloses, hydroxypropyl methylcellulose, carboxymethylcellulose sodium, methylcellulose, hydroxyethylcellulose, hydroxypropylmethylcellulose phthalate, noncrystalline cellulose, magnesium aluminum silicate, triethanolamine, polyvinyl alcohol, polyvinylpyrrolidone, 4-(1,1,3,3-tetramethylbutyl)-phenol polymer with ethylene oxide and formaldehyde, poloxamers; poloxamines, a charged phospholipid, dioctylsulfosuccinate, dialkylesters of sodium sulfosuccinic acid, sodium lauryl sulfate, alkyl aryl polyether sulfonates, mixtures of sucrose stearate and sucrose distearate,

$C_{18}H_{37}CH_2C(O)N(CH_3)-CH_2(CHOH)_4(CH_2OH)_2$, p-isononylphenoxypoly-(glycidol), decanoyl-N-methylglucamide; n-decyl β -D-glucopyranoside; n-decyl β -D-maltopyranoside; n-dodecyl β -D-glucopyranoside; n-dodecyl β -D-maltoside; heptanoyl-N-methylglucamide; n-heptyl- β -D-glucopyranoside; n-heptyl β -D-thioglucoside; n-hexyl β -D-glucopyranoside; nonanoyl-N-methylglucamide; n-noyl β -D-glucopyranoside; octanoyl-N-methylglucamide; n-octyl- β -D-glucopyranoside; octyl β -D-thioglucopyranoside; lysozyme, PEG-phospholipid, PEG-cholesterol, PEG-cholesterol derivative, PEG-vitamin A, PEG-vitamin E, and random copolymers of vinyl acetate and vinyl pyrrolidone.

61. (Previously Presented) The method of claim 59, wherein at least one cationic surface stabilizer is selected from the group consisting of a polymer, a biopolymer, a polysaccharide, a cellulosic, an alginate, a nonpolymeric compound, and a phospholipid.

62. (Currently Amended) The method of claim 59, wherein the surface stabilizer is selected from the group consisting of benzalkonium chloride, polymethylmethacrylate trimethylammonium bromide, polyvinylpyrrolidone-2-dimethylaminoethyl methacrylate dimethyl sulfate, hexadecyltrimethyl ammonium bromide, cationic lipids, sulfonium compounds, phosphonium compounds, quaternary ammonium compounds, benzyl-di(2-chloroethyl)ethylammonium bromide, coconut trimethyl ammonium chloride, coconut trimethyl ammonium bromide, coconut methyl dihydroxyethyl ammonium chloride, coconut methyl dihydroxyethyl ammonium bromide, decyl triethyl ammonium chloride, decyl dimethyl hydroxyethyl ammonium chloride, decyl dimethyl hydroxyethyl ammonium chloride bromide, C_{12-15} dimethyl hydroxyethyl ammonium chloride, C_{12-15} dimethyl hydroxyethyl ammonium chloride bromide, coconut dimethyl hydroxyethyl ammonium chloride, coconut dimethyl hydroxyethyl ammonium bromide, myristyl trimethyl ammonium methyl sulphate, lauryl dimethyl benzyl ammonium chloride, lauryl dimethyl benzyl ammonium bromide, lauryl dimethyl (ethenoxy)₄ ammonium chloride, lauryl dimethyl (ethenoxy)₄ ammonium bromide, N-alkyl (C_{12-18})dimethylbenzyl ammonium chloride, N-alkyl (C_{14-18})dimethyl-benzyl ammonium chloride, N-tetradecylidmethylbenzyl ammonium chloride monohydrate, dimethyl didecyl

ammonium chloride, N-alkyl and (C₁₂₋₁₄) dimethyl 1-naphthylmethyl ammonium chloride, trimethylammonium halide, alkyl-trimethylammonium salts, dialkyl-dimethylammonium salts, lauryl trimethyl ammonium chloride, ethoxylated alkyamidoalkyldialkylammonium salt, an ethoxylated trialkyl ammonium salt, dialkylbenzene dialkylammonium chloride, N-didecyldimethyl ammonium chloride, N-tetradecyldimethylbenzyl ammonium, chloride monohydrate, N-alkyl(C₁₂₋₁₄) dimethyl 1-naphthylmethyl ammonium chloride, dodecyldimethylbenzyl ammonium chloride, dialkyl benzenealkyl ammonium chloride, lauryl trimethyl ammonium chloride, alkylbenzyl methyl ammonium chloride, alkyl benzyl dimethyl ammonium bromide, C₁₂ trimethyl ammonium bromides, C₁₅ trimethyl ammonium bromides, C₁₇ trimethyl ammonium bromides, dodecylbenzyl triethyl ammonium chloride, poly-diallyldimethylammonium chloride (DADMAC), dimethyl ammonium chlorides, alkyldimethylammonium halogenides, tricetyl methyl ammonium chloride, decyltrimethylammonium bromide, dodecyltriethylammonium bromide, tetradecyltrimethylammonium bromide, methyl trioctylammonium chloride, ~~POLYQUAT 10™~~ polyquaternium 10, tetrabutylammonium bromide, benzyl trimethylammonium bromide, choline esters, benzalkonium chloride, stearylalkonium chloride compounds, cetyl pyridinium bromide, cetyl pyridinium chloride, halide salts of quaternized polyoxyethylalkylamines, ~~MIRAPOL™~~, ~~ALKAQUAT™~~ quaternized ammonium salt polymers, alkyl pyridinium salts; amines, amine salts, amine oxides, imide azolinium salts, protonated quaternary acrylamides, methylated quaternary polymers, and cationic guar.

63. (Previously Presented) The method of claim 50, wherein the T_{max} of the nanoparticulate composition, when assayed in the plasma of the mammalian subject, is selected from the group consisting of less than about 5 hours, less than about 4 hours, less than about 3 hours, less than 2 hours, less than about 1 hour, less than less than about 50 minutes, less than about 45 minutes, less than about 40 minutes, less than about 35 minutes, less than about 30 minutes, less than about 20 minutes, less than about 15 minutes, less than about 10 minutes, and less than about 5 minutes.

64. (Previously Presented) The method of claim 50, wherein the C_{max} of the nanoparticulate composition, when assayed in the plasma of the mammalian subject, is selected from the group consisting of greater than about 1 $\mu\text{g/mL}$, greater than about 3 $\mu\text{g/mL}$, greater than about 5 $\mu\text{g/mL}$, greater than about 10 $\mu\text{g/mL}$, and greater than about 15 $\mu\text{g/mL}$.

65. (Previously Presented) The method of claim 50, wherein the method is used to treat a condition selected from the group consisting of conditions in which NSAIDs are contraindicated, arthritic disorders, gastrointestinal conditions, inflammatory conditions, pulmonary inflammation, ophthalmic diseases, central nervous systems disorders, pain, fever, inflammation-related cardiovascular disorders, angiogenesis-related disorders, benign tumors, malignant tumors, adenomatous polyps, endometriosis, osteoporosis, dysmenorrhea, premature labor, asthma, fibrosis which occurs with radiation treatment, eosinophil-related disorders, pyrexia, bone resorption, nephrotoxicity, hypotension, arthrosis, joint stiffness, kidney disease, liver disease, acute mastitis, diarrhea, colonic adenomas, bronchitis, allergic neuritis, cytomegalovirus infectivity, apoptosis, lumbago, psoriasis, eczema, acne, burns, dermatitis, ultraviolet radiation damage, allergic rhinitis, respiratory distress syndrome, and endotoxin shock syndrome.

66. (Previously Presented) The method of claim 50, wherein the method is used to treat an indication in which anti-inflammatory agents, anti-angiogenesis agents, antitumorigenic agents, immunosuppressive agents, NSAIDs, COX-2 inhibitors, analgesic agents, anti-thrombotic agents, narcotics, or antifebrile agents are typically used.

67. (Previously Presented) The method of claim 50, wherein said subject is a human.

68. (Previously Presented) The method of claim 50, wherein said composition additionally comprising at least one non-meloxicam active agent selected from the group consisting of proteins, peptides, nucleotides, anti-obesity drugs, nutraceuticals, dietary supplements, carotenoids, corticosteroids, elastase inhibitors, anti-fungals, alkylxanthine, oncology therapies, anti-emetics, analgesics, opioids, antipyretics, cardiovascular agents, anti-

inflammatory agents, anthelmintics, anti-arrhythmic agents, antibiotics, anticoagulants, antidepressants, antidiabetic agents, antiepileptics, antihistamines, antihypertensive agents, antimuscarinic agents, antimycobacterial agents, antineoplastic agents, immunosuppressants, antithyroid agents, antiviral agents, anxiolytics, sedatives, astringents, beta-adrenoceptor blocking agents, blood products, blood substitutes, cardiac inotropic agents, contrast media, corticosteroids, cough suppressants, diagnostic agents, diagnostic imaging agents, diuretics, dopaminergics, haemostatics, immunological agents, lipid regulating agents, muscle relaxants, parasympathomimetics, parathyroid calcitonin and biphosphonates, prostaglandins, radio-pharmaceuticals, sex hormones, anti-allergic agents, stimulants, anoretics, sympathomimetics, thyroid agents, Mu receptor antagonists, Kappa receptor antagonists, non-narcotic analgesics, monoamine uptake inhibitors, adenosine regulating agents, cannabinoid derivatives, Substance P antagonists, neurokinin-1 receptor antagonists, and sodium channel blockers.

69. (Previously Presented) The method of claim 68, wherein said nutraceutical is selected from the group consisting of lutein, folic acid, fatty acids, fruit extracts, vegetable extracts, vitamin supplements, mineral supplements, phosphatidylserine, lipoic acid, melatonin, glucosamine/chondroitin, Aloe Vera, Guggul, glutamine, amino acids, green tea, lycopene, whole foods, food additives, herbs, phytonutrients, antioxidants, flavonoid constituents of fruits, evening primrose oil, flax seeds, fish oils, marine animal oils, and probiotics.

70. (Previously Presented) The method of claim 68, wherein said anti-inflammatory agent is a COX-2 inhibitor selected from the group consisting of celecoxib, rofecoxib, valdecoxib, parecoxib, MK-966, etoricoxib, 4-[5-(4-chlorophenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl]] benzenesulfonamide, N-(2-cyclohexyloxy-4-nitrophenyl)methane sulfonamide, methyl sulfone spiro(2.4)hept-5-ene I, SC-57666, celecoxib, SC-558, SC-560, etodolac, 5,5-dimethyl-3-(3-fluorophenyl)-4-(4-methylsulfonyl)phenyl 2(5H)-furanone, MK-476, L-745337, L-761066, L-761000, L-748780, L-748731, 5-Bromo-2-(4-fluorophenyl)-3-(4-(methylsulfonyl)phenyl, 1-(7-tert.-butyl-2,3-dihydro-3,3-dimethylbenzo(b)furan-5-yl)-4-cyclopropylbutan-1-one, 3-formylamino-7-methylsulfonylamino-6-phenoxy-4H-1-benzopyran-

4-one, BF 389, PD 136005, PD 142893, PD 145065, flurbiprofen, nimesulide, nabumetone, flosulide, piroxicam, diclofenac, COX-189, D 1367, 4 nitro 2 phenoxy methane sulfonanilide, (3 benzoyldifluoromethane sulfonanilide, diflumidone), JTE-522, 4'-Acetyl-2'-(2,4-difluorophenoxy)methanesulfonanilide, FK 867, FR 115068, GR 253035, RWJ 63556, RWJ 20485, ZK 38997, (E)-(5)-(3,5-di-tert-butyl-4-hydroxybenzylidene)-2-ethyl-1,2-isothiazolidine-1,1-dioxide indomethacin, CL 1004, RS 57067, RS 104894, SC 41930, SB 205312, SKB 209670, and Ono 1078.

71. (Previously Presented) The method of claim 68, wherein said non-meloxicam active agent is selected from the group consisting of aceclofenac, acetaminophen, acetaminosalol, acetanilide, acetylsalicylic acid, S-adenosylmethionine, alclofenac, alfentanil, allylprodine, alminoprofen, aloxiprin, alphaprodine, aluminum bis(acetylsalicylate), amfenac, aminochlorthenoxazin, 3-amino-4-hydroxybutyric acid, 2-amino-4-picoline, aminopropyl, aminopyrine, amixetrine, ammonium salicylate, ampiroxicam, amtolmetin guacil, anileridine, antipyrine, antipyrine salicylate, antrafenine, apazone, bendazac, benorylate, benoxaprofen, benzpiperylon, benzydamine, benzylmorphine, bermoprofen, bezitramide, α -bisabolol, bromfenac, p-bromoacetanilide, 5-bromosalicylic acid acetate, bromosaligenin, buccetin, buclocic acid, bucolome, bufexamac, bumadizon, buprenorphine, butacetin, butibufen, butophanol, calcium acetylsalicylate, carbamazepine, carbiphen, carprofen, carsalam, chlorobutanol, chlorthenoxazin, choline salicylate, cinchophen, cinmetacin, cirmadol, clidanac, clometacin, clonitazene, clonixin, clopirac, clove, codeine, codeine methyl bromide, codeine phosphate, codeine sulfate, cropropamide, crotethamide, desomorphine, dexoadrol, dextromoramide, dezocine, diampromide, diclofenac sodium, difenamizole, difenpiramide, diflunisal, dihydrocodeine, dihydrocodeinone enol acetate, dihydromorphine, dihydroxyaluminum acetylsalicylate, dimenoxadol, dimepheptanol, dimethylthiambutene, dioxaphetyl butyrate, dipipanone, diprocetyl, dipyrone, ditazol, droxicam, emorfazone, enfenamic acid, epirizole, eptazocine, etersalate, ethenzamide, ethoheptazine, ethoxazene, ethylmethylthiambutene, ethylmorphine, etodolac, etofenamate, etonitazene, eugenol, felbinac,

fenbufen, fenclozic acid, fendosal, fenoprofen, fentanyl, fentiazac, fepradinol, feprazone, floctafenine, flufenamic acid, flunoxaprofen, fluoresone, flupirtine, fluproquazone, flurbiprofen, fosfosal, gentisic acid, glafenine, glucametacin, glycol salicylate, guaiazulene, hydrocodone, hydromorphone, hydroxypethidine, ibufenac, ibuprofen, ibuproxam, imidazole salicylate, indomethacin, indoprofen, isofezolac, isoladol, isomethadone, isonixin, isoxepac, isoxicam, ketobemidone, ketoprofen, ketorolac, p-lactophenetide, lefetamine, levorphanol, lofentanil, lonazolac, lomoxicam, loxoprofen, lysine acetylsalicylate, magnesium acetylsalicylate, meclofenamic acid, mefenamic acid, meperidine, meptazinol, mesalamine, metazocine, methadone hydrochloride, methotrimепrazine, metiazinic acid, metofoline, metopon, mofebutazone, mofezolac, morazone, morphine, morphine hydrochloride, morphine sulfate, morpholine salicylate, myrophine, nabumetone, nalbuphine, 1-naphthyl salicylate, naproxen, narceine, nefopam, nicomorphine, nifenazone, niflumic acid, nimesulide, 5'-nitro-2'-propoxyacetanilide, norlevorphanol, normethadone, normorphine, norpipanone, olsalazine, opium, oxaceprol, oxametacine, oxaprozin, oxycodone, oxymorphone, oxyphenbutazone, papaveretum, paranyline, parsalimide, pentazocine, perisoxal, phenacetin, phenadoxone, phenazocine, phenazopyridine hydrochloride, phenocoll, phenoperidine, phenopyrazone, phenyl acetylsalicylate, phenylbutazone, phenyl salicylate, phenyramidol, piketoprofen, piminodine, pipebuzone, piperylone, piprofen, pirazolac, piritramide, piroxicam, pranoprofen, proglumetacin, proheptazine, promedol, propacetamol, propiram, propoxyphene, propyphenazone, proquazone, protizinic acid, ramifenazone, remifentanil, rimazolium metilsulfate, salacetamide, salicin, salicylamide, salicylamide o-acetic acid, salicylsulfuric acid, salsalte, salverine, simetride, sodium salicylate, sufentanil, sulfasalazine, sulindac, superoxide dismutase, suprofen, suxibuzone, talniflumate, tenidap, tenoxicam, terofenamate, tetrandrine, thiazolinobutazone, tiaprofenic acid, tiaramide, tilidine, tinoridine, tolfenamic acid, tolmetin, tramadol, tropesin, viminol, xenbucin, ximoprofen, zaltoprofen, and zomepirac.

72. (Previously Presented) The method of any of claims 68, 69, 70, or 71, wherein at least one non-meloxicam active agent has an effective average particle size of less than about 2 microns.

73. (Cancelled)